

TORELLI THEOREM FOR THE MODULI SPACES OF PAIRS

VICENTE MUÑOZ

ABSTRACT. Let X be a smooth projective curve of genus $g \geq 2$ over $\mathbb{C}$. A pair (E, ϕ) over X consists of an algebraic vector bundle E over X and a section $\phi \in H^0(E)$. There is a concept of stability for pairs which depends on a real parameter τ . Here we prove that the third cohomology groups of the moduli spaces of τ -stable pairs with fixed determinant and rank $n \geq 2$ are polarised pure Hodge structures, and they are isomorphic to $H^1(X)$ with its natural polarisation (except in very few exceptional cases). This implies a Torelli theorem for such moduli spaces. We recover that the third cohomology group of the moduli space of stable bundles of rank $n \geq 2$ and fixed determinant is a polarised pure Hodge structure, which is isomorphic to $H^1(X)$. We also prove Torelli theorems for the corresponding moduli spaces of pairs and bundles with non-fixed determinant.

1. INTRODUCTION

Let X be a smooth projective curve of genus $g \geq 2$ over the field of complex numbers. Fix $n \geq 2$ and $d \in \mathbb{Z}$. We shall denote by $M_X(n, L_0)$ the moduli space of polystable bundles E over X of rank n and determinant $\det(E) = L_0$, where L_0 is a line bundle of degree d . This is a projective variety, which is smooth over the locus of stable bundles. If n and d are coprime, then there are no properly semistable bundles, and $M_X(n, L_0)$ is smooth and projective. If n and d are not coprime, then the open subset of stable bundles $M_X^s(n, L_0) \subset M_X(n, L_0)$ is a smooth quasi-projective variety, and $M_X(n, L_0)$ is in general singular.

A pair (E, ϕ) over X consists of a bundle E of rank n and determinant $\det(E) = L_0$ over X together with a section $\phi \in H^0(E)$. There is a concept of stability for a pair which depends on the choice of a parameter $\tau \in \mathbb{R}$. This gives a collection of moduli spaces of τ -polystable pairs $\mathfrak{M}_X(\tau; n, L_0)$, which are projective varieties. It contains a smooth open subset $\mathfrak{M}_X^s(\tau; n, L_0) \subset \mathfrak{M}_X(\tau; n, L_0)$ consisting of τ -stable pairs. Pairs are discussed at length in [2, 4, 8, 12].

The range of the parameter τ is an open interval $I = I_{n,d} = (\tau_m, \tau_M) \subset \mathbb{R}$. This interval is split by a finite number of critical values τ_c . For a non-critical value $\tau \in I$, there are no properly semistable pairs, so $\mathfrak{M}_X(\tau; n, L_0) = \mathfrak{M}_X^s(\tau; n, L_0)$

2000 *Mathematics Subject Classification*. Primary: 14F45. Secondary: 14D20, 14H60.

Key words and phrases. Moduli space, complex curve, bundle, Torelli theorem.

Partially supported through grant MEC (Spain) MTM2007-63582.

is smooth and projective. For a critical value $\tau = \tau_c$, $\mathfrak{M}_X(\tau; n, L_0)$ is in general singular at properly τ -semistable points.

Our first main result computes the third cohomology group of $\mathfrak{M}_X^s(\tau; n, L_0)$.

Theorem 1.1. *Let $n \geq 2$. For $n = 2$ let $\tau_L = \tau_M - 1$, otherwise set $\tau_L = \tau_M$, and let $\tau \in (\tau_m, \tau_L)$. Assume that we are not in one of the following “bad” cases:*

- $(n, g, d) = (3, 2, 2k)$, $k = 1, 2, 3$;
- $(n, g, d) = (2, 2, 5), (2, 3, 5), (2, 2, 6)$, $\tau = \tau_M - 2$;
- $(n, g, d) = (3, 2, 5)$, $\tau = 2$.

Then

- (1) $H^3(\mathfrak{M}_X^s(\tau; n, L_0))$ *is a pure Hodge structure which is naturally polarised.*
- (2) *There is an isomorphism $H^3(\mathfrak{M}_X^s(\tau; n, L_0)) \cong H^1(X)$ of polarised Hodge structures.*

For $n = 2$, the moduli space $\mathfrak{M}_X(\tau; n, L_0)$ for $\tau \in (\tau_L, \tau_M)$ is a projective space. Therefore the above result for the third cohomology does not hold. On the other hand, the special cases that we remove are of low genus, rank and degree, and for particular critical values of τ .

The following corollary is a Torelli theorem for the moduli spaces of τ -stable pairs.

Corollary 1.2. *Let X be a smooth projective curve, $n \geq 2$, L_0 a line bundle of degree d over X . For $n = 2$ let $\tau_L = \tau_M - 1$, otherwise set $\tau_L = \tau_M$, and let $\tau \in (\tau_m, \tau_L)$. Consider another collection X' , n' , L'_0 , d' and τ' . Assume that (n, g, d, τ) and (n', g', d', τ') are not one of the exceptional cases enumerated in Theorem 1.1. Then*

- *If $\mathfrak{M}_X(\tau; n, L_0)$ and $\mathfrak{M}_{X'}(\tau'; n', L'_0)$ are isomorphic algebraic varieties, then $X \cong X'$.*
- *If $\mathfrak{M}_X^s(\tau; n, L_0)$ and $\mathfrak{M}_{X'}^s(\tau'; n', L'_0)$ are isomorphic algebraic varieties, then $X \cong X'$.*

The first statement is reduced to the second one since $\mathfrak{M}_X^s(\tau; n, L_0)$ is the smooth locus of $\mathfrak{M}_X(\tau; n, L_0)$. The second one is proved by looking at the polarised Hodge structure $H^3(\mathfrak{M}_X^s(\tau; n, L_0)) \cong H^1(X)$ and recovering X from $H^1(X)$ via the usual Torelli theorem.

For values of τ slightly bigger than τ_m , there is a natural map $\mathfrak{M}_X(\tau; n, L_0) \rightarrow M_X(n, L_0)$. This allows to prove the following result.

Theorem 1.3. *Let $n \geq 2$. Assume that $(n, g, d) \neq (2, 2, 2k), (3, 2, 3k), (2, 3, 2k)$, $k \in \mathbb{Z}$. Then*

- (1) $H^3(M_X^s(n, L_0))$ *is a pure Hodge structure which is naturally polarised.*
- (2) *There is an isomorphism $H^3(M_X^s(n, L_0)) \cong H^1(X)$ of polarised Hodge structures.*

For $(n, g, d) = (2, 2, \text{even})$, the moduli space $M_X(n, L_0)$ is isomorphic to $\mathbb{P}^3$ (see [14]), hence the above result does not hold. Also, $M_X^s(2, L_0) = M_X(2, L_0) - S$, where $S = \text{Jac}^d X / \pm 1$. From this it is easy to see that the result does not hold for $M_X^s(2, L_0)$ either.

For $(n, g, d) = (2, 3, 2k), (3, 2, 3k), k \in \mathbb{Z}$, $H^3(M_X^s(n, L_0))$ is a mixed Hodge structure, whose graded piece $\text{Gr}_W^3 H^3(M_X^s(n, L_0))$ is isomorphic to $H^1(X)$. However we are not able to polarise it with the methods in this paper.

Corollary 1.4. *Let X be a curve, $(n, g, d) \neq (2, 2, 2k), (3, 2, 3k), (2, 3, 2k), k \in \mathbb{Z}$. If X' is another curve, $n' \geq 2$, and L'_0 is a line bundle over X' of degree d' , and $(n', g', d') \neq (2, 2, 2k), (3, 2, 3k), (2, 3, 2k), k \in \mathbb{Z}$, then a Torelli theorem holds:*

- *If $M_X(n, L_0)$ and $M_{X'}(n', L'_0)$ are isomorphic algebraic varieties, then $X \cong X'$.*
- *If $M_X^s(n, L_0)$ and $M_{X'}^s(n', L'_0)$ are isomorphic algebraic varieties, then $X \cong X'$.*

When n and d are coprime, Theorem 1.3 has been proved in [13, 17, 10]. In the non-coprime case, the Torelli theorem was proved in [9] by different methods. Theorem 1.3 has been proved by Arapura and Sastry [1], but under the condition $g > \frac{3}{n-1} + \frac{n^2+3n+1}{2}$. Here we remove this lower bound assumption.

Our strategy of proof is the following. First, we find more convenient to rephrase the problem in terms of triples. A triple (E_1, E_2, ϕ) consists of a pair of bundles E_1, E_2 of ranks n_1, n_2 , with $\det(E_1) = L_1, \det(E_2) = L_2$, respectively, over X and a homomorphism $\phi : E_2 \rightarrow E_1$. Here L_1, L_2 are fixed line bundles of degrees d_1, d_2 , respectively. There is a suitable concept of stability for triples depending on a real parameter σ . This gives rise to moduli spaces $\mathcal{N}_X(\sigma; n_1, n_2, L_1, L_2)$ of σ -polystable triples.

There is an identification of moduli spaces of pairs and triples given by

$$\mathfrak{M}_X(\tau; n, L_0) \rightarrow \mathcal{N}_X(\sigma; n, 1, L_0, \mathcal{O}), \quad (E, \phi) \mapsto (E, \mathcal{O}, \phi),$$

where $\mathcal{O}$ is the trivial line bundle, and $\sigma = (n+1)\tau - d$. Actually, this rephrasing is a matter of aesthetic. The arguments can be carried out directly with the moduli spaces of pairs, but the formulas which appear using triples look more symmetric, and clearly they could eventually be generalised to the case of triples of arbitrary ranks n_1, n_2 .

The range of the parameter σ is an interval $I = (\sigma_m, \sigma_M) \subset \mathbb{R}$ split by a finite number of critical values σ_c . When σ moves without crossing a critical value, then $\mathcal{N}_{X, \sigma} = \mathcal{N}_X(\sigma; n, 1, L_1, L_2)$ remains unchanged, but when σ crosses a critical value, $\mathcal{N}_{X, \sigma}$ undergoes a birational transformation which we call a *flip*. We compute the codimension of the locus where this birational map is not an isomorphism to be at least 2, except in the bad case $n = 2, \sigma = \sigma_M - 3$ (corresponding to

$\tau = \tau_M - 1$). This allows to prove that the Hodge structures $H^3(\mathcal{N}_{X,\sigma}^s)$ are identified for different values of $\sigma \in I$. An explicit description of the moduli space $\mathcal{N}_{X,\sigma_M^-}$, where $\sigma_M^- = \sigma_M - \epsilon$, $\epsilon > 0$ small, allows to compute $H^3(\mathcal{N}_{X,\sigma_M^-})$ by induction on n .

For $\sigma = \sigma_m^+ = \sigma_m + \epsilon$, $\epsilon > 0$ small, we have a morphism $\mathcal{N}_{X,\sigma_m^+} \rightarrow M_X(n, L_0)$, $L_0 = L_1 \otimes L_2^{-n}$. This is a fibration over the locus $M_X^s(n, L_0)$, when $d = \deg(L_0)$ is large enough. A computation of the codimension of the locus of strictly semistable bundles allows to check that $H^3(M_X^s(n, L_0)) \cong H^3(\mathcal{N}_{X,\sigma_m^+})$.

We end up with the study of the case of non-fixed determinant. Let $M_X(n, d)$ denote the moduli space of semistable bundles of rank n and degree d over X . The open subset consisting of stable bundles will be denoted $M_X^s(n, d) \subset M_X(n, d)$. There is a natural map $M_X(n, d) \rightarrow \text{Jac}^d X$, whose fiber over L_0 is $M_X(n, L_0)$. Also let $\mathfrak{M}_X(\tau; n, d)$ be the moduli space of τ -semistable pairs (E, ϕ) , where E is a bundle of rank n and degree d . There is a map $\mathfrak{M}_X(\tau; n, d) \rightarrow \text{Jac}^d X$ as before. Denote by $\mathfrak{M}_X^s(\tau; n, d)$ the open subset of τ -stable triples. The following theorem is proved by reducing to the case of fixed determinant.

Corollary 1.5. *The following Torelli theorems hold. Let X, X' be projective smooth curves of genus $g, g' \geq 2$. Let $n, n' \geq 2$, and $d, d' \in \mathbb{Z}$.*

Let $\tau \in (\tau_m, \tau_L)$, $\tau' \in (\tau'_m, \tau'_L)$. Assume that (n, g, d, τ) and (n', g', d', τ') are not in one of the bad cases enumerated in Theorem 1.1. Then

- *If $\mathfrak{M}_X(\tau; n, d) \cong \mathfrak{M}_{X'}(\tau'; n', d')$ then $X \cong X'$.*
- *If $\mathfrak{M}_X^s(\tau; n, d) \cong \mathfrak{M}_{X'}^s(\tau'; n', d')$ then $X \cong X'$.*

Assume that $(n, g, d), (n'g', d')$ are not of the form $(2, 2, 2k), (3, 2, 3k), (2, 3, 2k)$, $k \in \mathbb{Z}$. Then

- *If $M_X(n, d) \cong M_{X'}(n', d')$ then $X \cong X'$.*
- *If $M_X^s(n, d) \cong M_{X'}^s(n', d')$ then $X \cong X'$.*

2. MODULI SPACES OF TRIPLES

Let X be a smooth projective curve of genus $g \geq 2$ over $\mathbb{C}$. A triple $T = (E_1, E_2, \phi)$ on X consists of two vector bundles E_1 and E_2 over X , of ranks n_1 and n_2 and degrees d_1 and d_2 , respectively, and a homomorphism $\phi: E_2 \rightarrow E_1$. We shall refer to (n_1, n_2, d_1, d_2) as the *type* of the triple.

For any $\sigma \in \mathbb{R}$ the σ -slope of T is defined by

$$\mu_\sigma(T) = \frac{d_1 + d_2}{n_1 + n_2} + \sigma \frac{n_2}{n_1 + n_2}.$$

We say that a triple $T = (E_1, E_2, \phi)$ is σ -stable if $\mu_\sigma(T') < \mu_\sigma(T)$ for any proper subtriple $T' = (E'_1, E'_2, \phi')$. We define σ -semistability by replacing the above strict

inequality with a weak inequality. A triple T is σ -polystable if it is the direct sum of σ -stable triples of the same σ -slope. We denote by

$$\mathcal{N}_X(\sigma; n_1, n_2, d_1, d_2)$$

the moduli space of σ -polystable triples of type (n_1, n_2, d_1, d_2) . This moduli space was constructed in [5] and [15]. It is a complex projective variety. The open subset of σ -stable triples will be denoted by $\mathcal{N}_X^s(\sigma; n_1, n_2, d_1, d_2)$.

Let L_1, L_2 be two bundles of degrees d_1, d_2 respectively. Then the moduli spaces of σ -semistable triples $T = (E_1, E_2, \phi)$ with $\det(E_1) = L_1$ and $\det(E_2) = L_2$ will be denoted

$$\mathcal{N}_X(\sigma; n_1, n_2, L_1, L_2),$$

and $\mathcal{N}_X^s(\sigma; n_1, n_2, L_1, L_2)$ will be the open subset of σ -stable triples.

Let $\mu(E) = \deg(E)/\text{rk}(E)$ denote the slope of a bundle E , and let $\mu_i = \mu(E_i) = d_i/n_i$, for $i = 1, 2$. Write

$$\begin{aligned} \sigma_m &= \mu_1 - \mu_2, \\ \sigma_M &= \begin{cases} \left(1 + \frac{n_1+n_2}{|n_1-n_2|}\right) (\mu_1 - \mu_2), & \text{if } n_1 \neq n_2, \\ \infty, & \text{if } n_1 = n_2, \end{cases} \end{aligned}$$

and let I denote the interval $I = (\sigma_m, \sigma_M)$. Then a necessary condition for $\mathcal{N}_X^s(\sigma; n_1, n_2, d_1, d_2)$ to be non-empty is that $\sigma \in I$ (see [6]). Note that $\sigma_m > 0$. To study the dependence of the moduli spaces on the parameter σ , we need to introduce the concept of critical value [5, 12].

Definition 2.1. The values of $\sigma_c \in I$ for which there exist $0 \leq n'_1 \leq n_1$, $0 \leq n'_2 \leq n_2$, d'_1 and d'_2 , with $n'_1 n_2 \neq n_1 n'_2$, such that

$$\sigma_c = \frac{(n_1 + n_2)(d'_1 + d'_2) - (n'_1 + n'_2)(d_1 + d_2)}{n'_1 n_2 - n_1 n'_2}, \quad (2.1)$$

are called *critical values*.

The interval I is split by a finite number of values $\sigma_c \in I$. The stability and semistability criteria for two values of σ lying between two consecutive critical values are equivalent; thus the corresponding moduli spaces are isomorphic. When σ crosses a critical value, the moduli space undergoes a transformation which we call a *flip*. We shall study the flips in some detail in the next section.

Relationship with pairs. A pair (E, ϕ) over X consists of a vector bundle E of rank n and with $\det(E) = L_0$, where L_0 is some fixed bundle of degree d , and $\phi \in H^0(E)$. Let $\tau \in \mathbb{R}$. We say that (E, ϕ) is τ -stable (see [8, Definition 4.7]) if:

- For any subbundle $E' \subset E$, we have $\mu(E') < \tau$.
- For any subbundle $E' \subset E$ with $\phi \in H^0(E')$, we have $\mu(E/E') > \tau$.

The concept of τ -semistability is defined by replacing the strict inequalities by weak inequalities. A pair (E, ϕ) is τ -polystable if $E = E' \oplus E''$, where $\phi \in H^0(E')$ and E'' is a polystable bundle of slope τ . The moduli space of τ -polystable pairs is denoted by $\mathfrak{M}_X(\tau; n, L_0)$.

Interpreting $\phi \in H^0(E)$ as a morphism $\phi : \mathcal{O} \rightarrow E$, where $\mathcal{O}$ is the trivial line bundle on X , we have a map $(E, \phi) \mapsto (E, \mathcal{O}, \phi)$ from pairs to triples. The τ -stability of (E, ϕ) corresponds to the σ -stability of $(E, \mathcal{O}, \phi)$, where (see [5])

$$\sigma = (n+1)\tau - d. \quad (2.2)$$

Therefore we have an isomorphism of moduli spaces

$$\mathfrak{M}_X(\tau; n, L_0) \cong \mathcal{N}_X(\sigma; n, 1, L_0, \mathcal{O}), \quad (2.3)$$

Alternatively, (2.3) may be taken as the definition of the moduli space of pairs. Note that σ_m and σ_M correspond under (2.2) to

$$\begin{aligned} \tau_m &= \frac{d}{n}, \\ \tau_M &= \frac{d}{n-1}. \end{aligned}$$

Theorem 2.2. *For non-critical values $\sigma \in I$, $\mathcal{N}_{X,\sigma} = \mathcal{N}_X(\sigma; n, 1, L_1, L_2)$ is smooth and projective, and it only consists of σ -stable points (i.e. $\mathcal{N}_{X,\sigma} = \mathcal{N}_{X,\sigma}^s$). For critical values $\sigma = \sigma_c$, $\mathcal{N}_{X,\sigma}$ is projective, and the open subset $\mathcal{N}_{X,\sigma}^s \subset \mathcal{N}_{X,\sigma}$ is smooth. The dimension of $\mathcal{N}_{X,\sigma}$ is $(n^2 - n - 1)(g - 1) + d_1 - nd_2 - 1$.*

Proof. In general, if σ is not a critical value for triples of type (n_1, n_2, d_1, d_2) and $\gcd(n_1, n_2, d_1 + d_2) = 1$, then σ -semistability is equivalent to σ -stability. This follows from [12, Remark 3.8].

Smoothness for the σ -stable points follows from [5, Proposition 6.3], since any σ -stable triple $T = (E_1, E_2, \phi)$ of type $(n, 1, d_1, d_2)$ satisfies automatically that $\phi : E_2 \rightarrow E_1$ is injective. The result is for non-fixed determinant, but the proof carries over to the case of fixed determinant.

The dimension appears in [8, Theorem 5.13] in the case of non-fixed determinant. Going over the proof, we see that we only have to subtract $2g$ to the formula in [8, Theorem 5.13]. $\square$

There is an isomorphism

$$\mathcal{N}_X(\sigma; n, 1, L_1, L_2) \cong \mathcal{N}_X(\sigma; n, 1, L_1 \otimes (L_2^*)^{\otimes n}, \mathcal{O}), \quad (2.4)$$

given by $(E_1, L_2, \phi) \mapsto (E_1 \otimes L_2^*, \mathcal{O}, \phi)$, so the moduli space (2.3) is as general as the moduli spaces $\mathcal{N}_X(\sigma; n, 1, L_1, L_2)$.

3. FLIPS FOR THE MODULI SPACES OF PAIRS

The homological algebra of triples is controlled by the hypercohomology of a certain complex of sheaves which appears when studying infinitesimal deformations [6, Section 3]. Let $T' = (E'_1, E'_2, \phi')$ and $T'' = (E''_1, E''_2, \phi'')$ be two triples of types (n'_1, n'_2, d'_1, d'_2) and $(n''_1, n''_2, d''_1, d''_2)$, respectively. Let $\text{Hom}(T'', T')$ denote the linear space of homomorphisms from T'' to T' , and let $\text{Ext}^1(T'', T')$ denote the linear space of equivalence classes of extensions of the form

$$0 \longrightarrow T' \longrightarrow T \longrightarrow T'' \longrightarrow 0,$$

where by this we mean a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & E'_1 & \longrightarrow & E_1 & \longrightarrow & E''_1 \longrightarrow 0 \\ & & \phi' \uparrow & & \phi \uparrow & & \phi'' \uparrow \\ 0 & \longrightarrow & E'_2 & \longrightarrow & E_2 & \longrightarrow & E''_2 \longrightarrow 0. \end{array}$$

To analyze $\text{Ext}^1(T'', T')$ one considers the complex of sheaves

$$C^\bullet(T'', T') : (E_1^{''*} \otimes E'_1) \oplus (E_2^{''*} \otimes E'_2) \xrightarrow{c} E_2^{''*} \otimes E'_1, \quad (3.1)$$

where the map c is defined by

$$c(\psi_1, \psi_2) = \phi' \psi_2 - \psi_1 \phi''.$$

We introduce the following notation:

$$\begin{aligned} \mathbb{H}^i(T'', T') &= \mathbb{H}^i(C^\bullet(T'', T')), \\ h^i(T'', T') &= \dim \mathbb{H}^i(T'', T'), \\ \chi(T'', T') &= h^0(T'', T') - h^1(T'', T') + h^2(T'', T'). \end{aligned}$$

By [6, Proposition 3.1], there are natural isomorphisms

$$\begin{aligned} \text{Hom}(T'', T') &\cong \mathbb{H}^0(T'', T'), \\ \text{Ext}^1(T'', T') &\cong \mathbb{H}^1(T'', T'). \end{aligned}$$

We shall use the following results later:

Lemma 3.1 ([11, Lemma 3.10]). *If $T'' = (E''_1, E''_2, \phi'')$ is an injective triple, that is $\phi'' : E''_2 \rightarrow E''_1$ is injective, then $\mathbb{H}^2(T'', T') = 0$.*

Proposition 3.2 ([6, Proposition 3.2]). *For any holomorphic triples T' and T'' we have*

$$\chi(T'', T') = (1 - g)(n''_1 n'_1 + n''_2 n'_2 - n''_2 n'_1) + n''_1 d'_1 - n'_1 d''_1 + n''_2 d'_2 - n'_2 d''_2 - n''_2 d'_1 + n'_1 d''_2.$$

Fix the type (n_1, n_2, d_1, d_2) for the moduli spaces of triples. For brevity, write $\mathcal{N}_\sigma = \mathcal{N}_X(\sigma; n_1, n_2, L_1, L_2)$. Let $\sigma_c \in I$ be a critical value and set

$$\sigma_c^+ = \sigma_c + \epsilon, \quad \sigma_c^- = \sigma_c - \epsilon,$$

where $\epsilon > 0$ is small enough so that σ_c is the only critical value in the interval (σ_c^-, σ_c^+) .

Definition 3.3. We define the *flip loci* as

$$\begin{aligned} \mathcal{S}_{\sigma_c^+} &= \{T \in \mathcal{N}_{\sigma_c^+}^s ; T \text{ is } \sigma_c^- \text{-unstable}\} \subset \mathcal{N}_{\sigma_c^+}^s, \\ \mathcal{S}_{\sigma_c^-} &= \{T \in \mathcal{N}_{\sigma_c^-}^s ; T \text{ is } \sigma_c^+ \text{-unstable}\} \subset \mathcal{N}_{\sigma_c^-}^s. \end{aligned}$$

It follows that (see [6, Lemma 5.3])

$$\mathcal{N}_{\sigma_c^+}^s - \mathcal{S}_{\sigma_c^+} = \mathcal{N}_{\sigma_c}^s = \mathcal{N}_{\sigma_c^-}^s - \mathcal{S}_{\sigma_c^-}.$$

Definition 3.4. Let $\sigma_c \in I$ be a critical value given by (n'_1, n'_2, d'_1, d'_2) in (2.1), and let $(n''_1, n''_2, d''_1, d''_2) = (n_1 - n'_1, n_2 - n'_2, d_1 - d'_1, d_2 - d'_2)$.

- (1) Define $\tilde{\mathcal{S}}_{\sigma_c^+}^0(n'_1, n'_2, d'_1, d'_2)$ to be the set of all isomorphism classes of extensions

$$0 \longrightarrow T' \longrightarrow T \longrightarrow T'' \longrightarrow 0,$$

where T' and T'' are σ_c^+ -stable triples with types (n'_1, n'_2, d'_1, d'_2) and $(n''_1, n''_2, d''_1, d''_2)$ respectively, and for which T is σ_c^+ -stable, $T \in \mathcal{N}_{\sigma_c^+}^s$. Note that in this case

$$\mu_{\sigma_c}(T') = \mu_{\sigma_c}(T) = \mu_{\sigma_c}(T''), \text{ and } \frac{n'_2}{n'_1 + n'_2} < \frac{n''_2}{n''_1 + n''_2}.$$

- (2) Define

$$\tilde{\mathcal{S}}_{\sigma_c^+}^0 = \bigcup \tilde{\mathcal{S}}_{\sigma_c^+}^0(n'_1, n'_2, d'_1, d'_2),$$

where the union is over all (n'_1, n'_2, d'_1, d'_2) and $(n''_1, n''_2, d''_1, d''_2)$ such that the above conditions apply.

- (3) Similarly, define $\tilde{\mathcal{S}}_{\sigma_c^-}^0(n'_1, n'_2, d'_1, d'_2)$ and $\tilde{\mathcal{S}}_{\sigma_c^-}^0$, where now $\frac{n'_2}{n'_1 + n'_2} > \frac{n''_2}{n''_1 + n''_2}$.

The following is [6, Lemma 5.8]. Actually the version in [6] is stated for the case of non-fixed determinant, but the fixed determinant version is completely similar.

Lemma 3.5. *There are maps $v^\pm : \tilde{\mathcal{S}}_{\sigma_c^\pm}^0 \longrightarrow \mathcal{N}_{\sigma_c^\pm}^s$ which map triples to their equivalence classes. The images contain the flip loci $\mathcal{S}_{\sigma_c^\pm}$.*

Proposition 3.6. *Assume that $\mathbb{H}^0(T'', T') = \mathbb{H}^2(T'', T') = 0$ for all $\sigma_c^\pm$ -stable triples T', T'' of types (n'_1, n'_2, d'_1, d'_2) , $(n''_1, n''_2, d''_1, d''_2)$ respectively. Then $\mathcal{S}_{\sigma_c^\pm} \subset \mathcal{N}_{\sigma_c^\pm}^s$ are contained in subvarieties of codimension bounded below by*

$$\min\{-\chi(T', T'')\},$$

where the minimum is over all (n'_1, n'_2, d'_1, d'_2) which satisfy (2.1), $(n''_1, n''_2, d''_1, d''_2) = (n_1 - n'_1, n_2 - n'_2, d_1 - d'_1, d_2 - d'_2)$ and $\frac{n'_2}{n'_1 + n'_2} < \frac{n''_2}{n''_1 + n''_2}$ (in the case of $\mathcal{S}_{\sigma_c^+}$) or $\frac{n'_2}{n'_1 + n'_2} > \frac{n''_2}{n''_1 + n''_2}$ (in the case of $\mathcal{S}_{\sigma_c^-}$).

Proof. The proof of Proposition 5.10 in [6] goes over to this case. The condition $\sigma_c > 2g - 2$ in [6, Proposition 5.10] is only needed to conclude the vanishing of $\mathbb{H}^0(T'', T')$ and $\mathbb{H}^2(T'', T')$.

Fixing the determinant of the triples T forces that, once T' is chosen, then the determinant of T'' is fixed. This reduces by $2g$ the dimension of the moduli space of $\sigma^\pm$ -stable triples, and also reduces by $2g$ the dimension of the flip loci. Therefore the formula of the codimension is the same as in [6, Proposition 5.10]. $\square$

4. CODIMENSION ESTIMATES

We are going to apply Proposition 3.6 to the case of $n_2 = 1$, $n_1 = n \geq 2$. Denote $\mathcal{N}_{X,\sigma} = \mathcal{N}_X(\sigma; n, 1, L_1, L_2)$. Here L_1, L_2 are line bundles of degrees d_1, d_2 respectively. (We shall have no need of particularising $L_2 = \mathcal{O}$ for the subsequent arguments to work, so we will not do it).

We start by computing codimension estimates for the flip loci $\mathcal{S}_{\sigma_c^\pm} \subset \mathcal{N}_{X,\sigma_c^\pm} = \mathcal{N}_{X,\sigma_c^\pm}^s$.

Proposition 4.1. *Suppose $n_2 = 1$, $n_1 = n \geq 2$. Let σ_c be a critical value with $\sigma_m < \sigma_c < \sigma_M$. Then*

- $\text{codim } \mathcal{S}_{\sigma_c^+} \geq 3$, except in the case $n = 2$, $g = 2$, d_1 odd and $\sigma_c = \sigma_m + \frac{3}{2}$ (in which case $\text{codim } \mathcal{S}_{\sigma_c^+} = 2$).
- $\text{codim } \mathcal{S}_{\sigma_c^-} \geq 2$, except for $n = 2$ and $\sigma_c = \sigma_M - 3$ (in which case $\text{codim } \mathcal{S}_{\sigma_c^-} = 1$). Moreover, for $n = 2$ we have that $\text{codim } \mathcal{S}_{\sigma_c^-} = 2$ only for $\sigma_c = \sigma_M - 6$.

Proof. Let us do the case of $\mathcal{S}_{\sigma_c^+}$ first. The condition

$$\frac{n'_2}{n'_1 + n'_2} < \frac{n''_2}{n''_1 + n''_2}$$

implies that $n'_2 = 0$ and $n''_2 = 1$. Since T' and T'' are σ_c^+ -stable triples which are not isomorphic, we have that $\mathbb{H}^0(T'', T') = \text{Hom}(T'', T') = 0$. By Lemma 3.1, it is clear that $\mathbb{H}^2(T'', T') = 0$. Proposition 3.2 gives (using that $n'_2 = d'_2 = 0$, $n''_2 = 1$, and paying attention to the fact that the roles of T' and T'' are interchanged)

$$-\chi(T', T'') = (g - 1)n'_1 n''_1 + n''_1 d'_1 - n'_1 d''_1.$$

The equality $\mu_{\sigma_c}(T') = \mu_{\sigma_c}(T)$ is rewritten as

$$\frac{d'_1}{n'_1} = \frac{d_1 + d_2 + \sigma_c}{n_1 + 1}.$$

Now $\sigma_c > \sigma_m = \frac{d_1}{n_1} - d_2$ implies that

$$\frac{d'_1}{n'_1} > \frac{1}{n_1 + 1} \left(d_1 + d_2 + \frac{d_1}{n_1} - d_2 \right) = \frac{d_1}{n_1}.$$

So $\frac{d'_1}{n'_1} > \frac{d''_1}{n''_1}$, and hence $n''_1 d'_1 - n'_1 d''_1 > 0$. This implies

$$-\chi(T', T'') \geq (g-1)n'_1 n''_1 + 1 \geq 2,$$

using that $n = n'_1 + n''_1$, $0 < n'_1, n''_1 < n$. Moreover, except in the case $(n, g) = (2, 2)$, we have that $-\chi(T', T'') \geq 3$. For $n = 2$, $g = 2$, $n'_1 = n''_1 = 1$, $-\chi(T', T'') = d'_1 - d''_1 + 1$, with $d'_1 > d''_1$. For $d'_1 - d''_1 = 1$, d_1 is odd, $d'_1 = (d_1 + 1)/2$ and $d''_1 = (d_1 - 1)/2$. Therefore $\sigma_c = (d_1 + 3)/2 - d_2 = \sigma_m + 3/2$.

Now we turn to the case of $\mathcal{S}_{\sigma_c^-}$. The condition

$$\frac{n'_2}{n'_1 + n'_2} > \frac{n''_2}{n''_1 + n''_2}$$

implies that $n'_2 = 1$ and $n''_2 = 0$. Since T' and T'' are σ_c^- -stable triples which are not isomorphic, we have that $\mathbb{H}^0(T'', T') = \text{Hom}(T'', T') = 0$. Lemma 3.1 guarantees that $\mathbb{H}^2(T'', T') = 0$. Proposition 3.2 gives (using that $n''_2 = d''_2 = 0$, $n'_2 = 1$)

$$-\chi(T', T'') = (g-1)n''_1(n'_1 - 1) + n''_1 d'_1 - n'_1 d''_1 + d''_1 - n''_1 d_2.$$

Denote

$$A = n''_1 d'_1 - n'_1 d''_1 + d''_1 - n''_1 d_2. \quad (4.1)$$

We have

$$A = n''_1(d_1 - d_2) - d''_1(n'_1 + n''_1 - 1) = n''_1(d_1 - d_2) - d''_1(n_1 - 1).$$

Also $\mu_{\sigma_c}(T'') = \mu_{\sigma_c}(T)$ means that

$$\frac{d''_1}{n''_1} = \frac{d_1 + d_2 + \sigma_c}{n_1 + 1},$$

From where

$$A = n''_1(d_1 - d_2) - n''_1(n_1 - 1) \frac{d_1 + d_2 + \sigma_c}{n_1 + 1}.$$

Now

$$\sigma_c < \sigma_M = \frac{2n_1}{n_1 - 1} \left(\frac{d_1}{n_1} - d_2 \right)$$

so that

$$A > n''_1(d_1 - d_2) - n''_1 \frac{n_1 - 1}{n_1 + 1} \left(d_1 + d_2 + \frac{2n_1}{n_1 - 1} \left(\frac{d_1}{n_1} - d_2 \right) \right) = 0.$$

This gives that

$$-\chi(T', T'') = (g-1)n''_1(n'_1 - 1) + A \geq 2,$$

in the case $n'_1 > 1$. In the case $n'_1 = 1$, we have a more explicit formula

$$-\chi(T', T'') = A = (n_1 - 1)(d'_1 - d_2) > 0.$$

Hence $-\chi(T', T'') \geq 2$, for $n \geq 3$ and for $n = 2$ and $d'_1 - d_2 \geq 2$. The only remaining case corresponds to $-\chi(T', T'') = d'_1 - d_2 = 1$, $n'_1 = n''_1 = 1$, $d'_1 = d_2 + 1$, $\sigma_M = 2d_1 - 4d_2$, $\sigma_m = d_1/2 - d_2$ and $\sigma_c = 3d''_1 - d_1 - d_2 = \sigma_M - 3$. Finally, note

that for $n = 2$, $-\chi(T', T'') = 2$ only in the case $d'_1 - d_2 = 2$, which corresponds to $\sigma_c = \sigma_M - 6$. $\square$

Lemma 4.2. *Let σ_c be a critical value. Assume that $\sigma_c < \sigma_L$ in the case $n = 2$. Then $\text{codim}(\mathcal{N}_{X, \sigma_c} - \mathcal{N}_{X, \sigma_c}^s) \geq 5$ except in the following cases:*

- $n = 2, g = 2, 3, \sigma_c = \sigma_M - 6$ and $d_1 - 2d_2 = 5$,
- $n = 2, g = 2, \sigma_c = \sigma_M - 6$ and $d_1 - 2d_2 = 6$,
- $n = 3, g = 2, \sigma_c = 2$ and $d_1 - 3d_2 = 4$,
- $n = 3, g = 2, \sigma_c = 3$ and $d_1 - 3d_2 = 5$.

Proof. By Theorem 2.2, the dimension of $\mathcal{N}_{X, \sigma}$ is

$$\dim \mathcal{N}_X(\sigma; n, 1, L_1, L_2) = (n^2 - n - 1)(g - 1) + d_1 - nd_2 - 1.$$

The set $S = \mathcal{N}_{X, \sigma_c} - \mathcal{N}_{X, \sigma_c}^s$ is formed by strictly σ_c -polystable triples. Therefore it is covered by the images of the sets

$$\mathcal{X}_{L_1, L_2} \subset \mathcal{N}_X^s(\sigma_c; n'_1, 1, d'_1, d_2) \times M_X(n''_1, d''_1),$$

where $d_1 = d'_1 + d''_1$, $n_1 = n'_1 + n''_1$,

$$\frac{d''_1}{n''_1} = \frac{d'_1 + d_2 + \sigma_c}{n'_1 + 1} = \frac{d_1 + d_2 + \sigma_c}{n_1 + 1}, \quad (4.2)$$

and $\mathcal{X}_{L_1, L_2}$ corresponds to those triples of the form $T = (E_1, L_2, \phi) = (E'_1, L_2, \phi) \oplus (E''_1, 0, 0)$ with fixed determinant $\det(E_1) = L_1$. Therefore $\mathcal{X}_{L_1, L_2}$ is a fibration over $M_X(n''_1, d''_1)$ whose fibers are moduli spaces of σ_c -stable triples with fixed determinant $\det(E'_1) = L_1 \otimes \det(E''_1)^{-1}$. Thus

$$\dim \mathcal{X} = (n_1'^2 - n'_1 - 1)(g - 1) + d'_1 - n'_1 d_2 - 1 + (n_1''^2(g - 1) + 1).$$

So

$$\begin{aligned} \text{codim } \mathcal{X} &= (n^2 - n + n'_1 - n_1'^2 - n_1''^2)(g - 1) + d_1 - d'_1 - (n - n'_1)d_2 - 1 \\ &= (2n'_1 n''_1 - n''_1)(g - 1) + d''_1 - n''_1 d_2 - 1 \\ &= n''_1(2n'_1 - 1)(g - 1) + d''_1 - n''_1 d_2 - 1. \end{aligned}$$

Note that

$$d''_1 - n''_1 d_2 = (n''_1 d'_1 - n'_1 d''_1 + d''_1 - n''_1 d_2) + (n'_1 d''_1 - n''_1 d'_1). \quad (4.3)$$

Define $A = n''_1 d'_1 - n'_1 d''_1 + d''_1 - n''_1 d_2$ as in (4.1). Using (4.2), we get as in the proof of Proposition 4.1 that $A > 0$. Also the inequality $\sigma_c > \frac{d'_1}{n'_1} - d_2 > 0$ and (4.2) give

$$\frac{d''_1}{n''_1} > \frac{d'_1 + d_2 + d'_1/n'_1 - d_2}{n'_1 + 1} = \frac{d'_1}{n'_1},$$

hence $B = n'_1 d''_1 - n''_1 d'_1 > 0$. To prove that $\text{codim } \mathcal{X} \geq 5$ we need to prove that

$$1 + \text{codim } \mathcal{X} = n''_1(2n'_1 - 1)(g - 1) + A + B \geq 6.$$

This is true except possibly in the following cases:

- $n = 2$. Then $n'_1 = 1$ and $n''_1 = 1$. We have that $A = d'_1 - d_2$ and $B = d''_1 - d'_1$. In this case we assume that $\sigma_c = 3d''_1 - d_1 - d_2 < \sigma_L = \sigma_M - 3 = 2d_1 - 4d_2 - 3$, which can be rewritten as $A = d'_1 - d_2 > 1$. So $1 + \text{codim } \mathcal{X} = (g - 1) + A + B \geq 6$ except for $g = 2, 3$, $A = 2$ and $B = 1$ (that is, $\sigma_c = \sigma_M - 6$ and $d_1 - 2d_2 = 5$), for $g = 2$, $A = 2$ and $B = 2$ (that is, $\sigma_c = \sigma_M - 6$ and $d_1 - 2d_2 = 6$).
- $n \geq 3$ and $n'_1 = 1$. Then $n''_1(2n'_1 - 1)(g - 1) = n''_1(g - 1) \geq 2$ and $A = n''_1(d'_1 - d_2) \geq 2$. So $1 + \text{codim } \mathcal{X} \geq 6$ except if $n''_1 = 2$, $g = 2$, $d'_1 - d_2 = 1$ and $B = d''_1 - n''_1 d'_1 = 1$. This means that $\sigma_c = 2d''_1 - d_1 - d_2 = 3d_2 - d_1 + 6$. As $\sigma_M = d_1 - 3d_2$, $\sigma_m = \frac{1}{3}(d_1 - 3d_2)$ and $\sigma_m < \sigma_c < \sigma_M$, we have that it must be $\sigma_c = 2$ and $d_1 - 3d_2 = 4$.
- $n \geq 3$ and $n'_1 \geq 2$. Then $n''_1(2n'_1 - 1)(g - 1) \geq 3$. So $1 + \text{codim } \mathcal{X} \geq 6$ except if $n''_1(2n'_1 - 1)(g - 1) = 3$, $A = 1$ and $B = 1$. This means $n'_1 = 2$, $n''_1 = 1$, $g = 2$, $A = d'_1 - d''_1 - d_2 = 1$, $B = 2d''_1 - d'_1 = 1$. So $\sigma_c = 4d''_1 - d_1 - d_2 = 3d_2 - d_1 + 8$. As $\sigma_M = d_1 - 3d_2$, $\sigma_m = \frac{1}{3}(d_1 - 3d_2)$ and $\sigma_m < \sigma_c < \sigma_M$, we have that it must be $\sigma_c = 3$ and $d_1 - 3d_2 = 5$.

□

Also we need codimension estimates for the families of properly semistable bundles over X .

Lemma 4.3. *Let S be a bounded family of isomorphism classes of strictly semistable bundles of rank n and determinant L_0 . Then $\dim M_X(n, L_0) - \dim S \geq (n - 1)(g - 1)$.*

Proof. We may stratify S according to the ranks and degrees of the elements in the Jordan-Hölder filtration of the bundles. So we may assume that S consists only of bundles whose Jordan-Hölder filtration has associated graded object $\text{gr}(Q) = \bigoplus_{i=1}^r Q_i$, with $n_i = \text{rk } Q_i$, $d_i = \text{deg } Q_i$, $\bigotimes_{i=1}^r \det(Q_i) = L_0$, $r \geq 2$. We use now [7, Proposition 7.9], but it has to be modified to take into account that we are fixing the determinant of Q . This fixes the determinant of one of the Q_i , say Q_r . This reduces the dimension stated in [7, Proposition 7.9] by g . So

$$\dim S \leq \left(\sum n_i^2 + \sum_{i < j} n_i n_j \right) (g - 1) + 1 - g.$$

As $\dim M_X(n, L_0) = n^2(g - 1) + 1 - g$, we have that

$$\dim M(n, L_0) - \dim S \geq \sum_{i < j} n_i n_j (g - 1).$$

The minimum of the right hand side is attained for $n = 2$, $n_1 = 1$, $n_2 = n - 1$, whence the statement. □

Lemma 4.4. *Suppose $n \geq 2$, and let $S = M_X(n, L_0) - M_X^s(n, L_0)$ be the locus of strictly polystable bundles. Then $\dim M_X(n, L_0) - \dim S \geq 2(n-1)(g-1) - 1$.*

Proof. Working as in the proof of Lemma 4.3, we have that if S consists of polystable bundles $Q = \oplus Q_i$, with $n_i = \text{rk } Q_i$, $d_i = \deg Q_i$, $\bigotimes_{i=1}^r \det(Q_i) = L_0$, $r \geq 2$, then

$$\dim S = \left(\sum n_i^2(g-1) + 1 \right) - g.$$

As $\dim M_X(n, L_0) = n^2(g-1) + 1 - g$, we have that

$$\dim M_X(n, L_0) - \dim S \geq 2 \sum_{i < j} n_i n_j (g-1) + 1 - r \geq 2(n-1)(g-1) - 1,$$

since the minimum occurs for $r = 2$, $n_1 = 1$, $n_2 = n - 1$. $\square$

5. COHOMOLOGY GROUPS OF THE MODULI SPACES OF PAIRS

Now we aim to compute the cohomology groups $H^i(\mathcal{N}_{X,\sigma}^s)$, for $i = 1, 2, 3$. Note that for σ non-critical, $\mathcal{N}_{X,\sigma} = \mathcal{N}_{X,\sigma}^s$, so we are actually computing $H^i(\mathcal{N}_{X,\sigma})$. Moreover, in this case $\mathcal{N}_{X,\sigma}$ is smooth and projective, so these are automatically pure Hodge structures. As a byproduct, we shall obtain the cohomology groups $H^i(M_X^s(n, L))$, for $i = 1, 2, 3$. For n and d coprime, $M_X(n, L) = M_X^s(n, L)$, which is smooth and projective. In this case, these cohomology groups are well-known, however we shall recover them easily from our arguments. For n and d not coprime, the cohomology groups $H^i(M_X^s(n, L))$ seem to be known to experts, but it is difficult to locate them in the literature.

We start with a small lemma.

Lemma 5.1. *Let M be a smooth projective variety and let $S \subset M$ be a closed subset in the Zariski topology. If either:*

- (1) $\text{codim } S \geq 3$, or
- (2) $\text{codim } S = 2$ and $H^3(M - S)$ is a pure Hodge structure,

then

$$H^i(M) \cong H^i(M - S),$$

for $i \leq 3$.

Proof. Let $m = \dim_{\mathbb{C}} M$. Using Poincaré duality, the statement of the lemma is equivalent to proving an isomorphism $H_c^{2m-i}(M) \cong H_c^{2m-i}(M - S)$, where H_c^* stands for cohomology with compact support. This is clear for $i \leq 2$, since S has real codimension at least 4. For $i = 3$, we have an exact sequence

$$H_c^{2m-4}(S) \xrightarrow{\partial} H_c^{2m-3}(M - S) \longrightarrow H_c^{2m-3}(M) \longrightarrow 0.$$

The group $H_c^{2m-4}(S)$ is generated by the irreducible components S_i of dimension $m - 2$ of S . To see this, let $N_1 = \bigcup_{i \neq j} (S_i \cap S_j)$. Then let $S_i^o \subset S_i$ be the smooth

locus of $S_i - N_1$, and consider $N_2 = \bigcup (S_i - S_i^o)$. Then $N = N_1 \cup N_2$ is of positive codimension in S , so that $H_c^{2m-4}(S) \cong H_c^{2m-4}(S - N) = \bigoplus_i H_c^{2m-4}(S_i^o)$. Note that $H_c^{2m-4}(S)$ is a pure Hodge structure of weight $(m-2, m-2)$. As ∂ preserves the weight of the Hodge structure, and $H_c^{2m-3}(M - S)$ is assumed to be of pure type, then $\partial = 0$. The result follows. $\square$

Remark 5.2. If $S \subset M$ is of codimension 2 and M is a smooth projective variety, then in general, $H^3(M - S)$ is a mixed Hodge structure. It has pieces of weight 2 and 3, and the piece of weight 3 satisfies $\mathrm{Gr}_3^W H^3(M - S) \cong H^3(M)$.

Proposition 5.3. Assume $n \geq 3$. Let σ_c be a critical value, $\sigma_m < \sigma_c < \sigma_M$. Then

$$H^i(\mathcal{N}_{X, \sigma_c^+}) \cong H^i(\mathcal{N}_{X, \sigma_c^-}) \cong H^i(\mathcal{N}_{X, \sigma_c}^s),$$

for $i \leq 3$. So all Hodge structures $H^i(\mathcal{N}_{X, \sigma}^s)$ are naturally isomorphic, for $i \leq 3$. (For σ non-critical, we have that $\mathcal{N}_{X, \sigma} = \mathcal{N}_{X, \sigma}^s$, so we are actually talking about $H^i(\mathcal{N}_{X, \sigma})$.)

Proof. By Proposition 4.1, $\mathrm{codim} \mathcal{S}_{\sigma_c^+} \geq 3$ and $\mathrm{codim} \mathcal{S}_{\sigma_c^-} \geq 2$. Then Lemma 5.1 applied to $\mathcal{S}_{\sigma_c^+}$ implies that $H^3(\mathcal{N}_{X, \sigma_c^+}) \cong H^3(\mathcal{N}_{X, \sigma_c}^s)$, since $\mathcal{N}_{X, \sigma}^s = \mathcal{N}_{X, \sigma_c^+} - \mathcal{S}_{\sigma_c^+}$. In particular, $H^3(\mathcal{N}_{X, \sigma_c}^s)$ is a pure Hodge structure. Applying Lemma 5.1 to $\mathcal{S}_{\sigma_c^-}$, we have $H^3(\mathcal{N}_{X, \sigma_c^-}) \cong H^3(\mathcal{N}_{X, \sigma_c}^s)$. $\square$

Proposition 5.4. Assume $n = 2$. The critical values are the numbers $\sigma_c = \sigma_M - 3n$, for $0 < n < (\sigma_M - \sigma_m)/3$, $n \in \mathbb{Z}$. Denote $\sigma_L = \sigma_M - 3$. If $\sigma_L > \sigma_m$ then for any $\sigma \in (\sigma_m, \sigma_L)$, we have $H^1(\mathcal{N}_{X, \sigma}^s) = 0$, $H^2(\mathcal{N}_{X, \sigma}^s) \cong \mathbb{Z} \oplus \mathbb{Z}$, and all Hodge structures $H^3(\mathcal{N}_{X, \sigma}^s)$ are naturally isomorphic. Moreover,

$$H^3(\mathcal{N}_{X, \sigma}^s) \cong H^1(X),$$

for any $\sigma \in (\sigma_m, \sigma_L)$, with the exception of $g = 2$, $d_1 - 2d_2 = 5$ and $\sigma = \sigma_M - 6 = 4$. (For σ non-critical, we have that $\mathcal{N}_{X, \sigma} = \mathcal{N}_{X, \sigma}^s$, so we are actually talking about $H^i(\mathcal{N}_{X, \sigma})$.)

Proof. The collection of moduli spaces $\mathcal{N}_{X, \sigma}$ for $n = 2$ is described in detail in [16]. The critical values are given by [12, Lemma 5.3] to be of the form $\sigma_c = 3(d_1 - d_2 - n) - d_1 - d_2 = \sigma_M - 3n$, with $n > 0$ and the constraint $\sigma_m < \sigma_c < \sigma_M$ (this also follows easily from (2.1)).

The last moduli space $\mathcal{N}_{X, \sigma_M^-}$ is a projective space $\mathbb{P}$ (see [16, (3.1)]), or argue as in the proof of Proposition 5.7 below: in the discussion of the proof of Proposition 5.7, F should be a fixed line bundle, since the determinant is fixed, so $\mathcal{N}_{X, \sigma_M^-} = U$ is the projective space $\mathbb{P} = \mathbb{P}H^1(F^* \otimes L)$.

By [16, (3.4)], there is an embedding $X \hookrightarrow \mathbb{P}$, given by $p \mapsto [\delta((F^* \otimes L(p))_p)] \in \mathbb{P}$, where $\delta : H^0((F^* \otimes L(p))_p) \rightarrow H^1(F^* \otimes L)$ is the connecting map associated to

the exact sequence

$$F^* \otimes L \rightarrow F^* \otimes L(p) \rightarrow (F^* \otimes L(p))_p.$$

By [16, (3.19)], the moduli space $\mathcal{N}_{X, \sigma_L^-}$, for $\sigma_L = \sigma_M - 3$, is the blow-up of $\mathbb{P}$ along X . The usual computation of the cohomology of a blow-up gives that $H^3(\mathcal{N}_{X, \sigma_L^-}) \cong H^1(E) \cong H^1(X)$, where E is the exceptional divisor, which is a projective bundle over X . Also $H^2(\mathcal{N}_{X, \sigma_L^-}) \cong H^2(\mathbb{P}) \oplus \mathbb{Z}[E] \cong \mathbb{Z} \oplus \mathbb{Z}$ and $H^1(\mathcal{N}_{X, \sigma_L^-}) = 0$.

Now we have to prove that

$$H^i(\mathcal{N}_{X, \sigma_c^+}) \cong H^i(\mathcal{N}_{X, \sigma_c^-}) \cong H^i(\mathcal{N}_{X, \sigma_c}^s),$$

for $i \leq 3$ and $\sigma_m < \sigma_c < \sigma_L$. In the case $g \neq 2$ (or in the case $g = 2$ and d_1 even or $\sigma_c \neq \sigma_m + \frac{3}{2}$), it follows from the fact that $\text{codim } \mathcal{S}_{\sigma_c^+} \geq 3$ and $\text{codim } \mathcal{S}_{\sigma_c^-} \geq 2$, which follows in turn from Proposition 4.1. This is enough to complete the proof of the proposition.

For the exceptional case $(n, g) = (2, 2)$, d_1 odd and $\sigma_c = \sigma_m + \frac{3}{2}$, we have the following cases:

- If $d_1 - 2d_2 = 1$ then there are no flips. So there is no such σ_c and nothing to prove. (Actually $\mathcal{N}_{X, \sigma}$ is a projective space for all allowable values of σ .)
- If $d_1 - 2d_2 = 3$ then $\sigma_c = \sigma_m + \frac{3}{2} = \sigma_M - 3$. Again there is no such σ_c . (Note that $\mathcal{N}_{X, \sigma_m^+} = \mathcal{N}_{X, \sigma_L^-}$, so for all $\sigma \in (\sigma_m, \sigma_L)$ we have $\mathcal{N}_{X, \sigma} = \mathcal{N}_{X, \sigma_m^+}$ whose cohomology has been computed above.)
- If $d_1 - 2d_2 \geq 7$ then $\sigma_m + \frac{3}{2} < \sigma_M - 6$. Then for $\sigma_c = \sigma_m + \frac{3}{2}$, $\text{codim } \mathcal{S}_{\sigma_c^+} = 2$ but $\text{codim } \mathcal{S}_{\sigma_c^-} \geq 3$ (see the last line in Proposition 3.6). Then $H^3(\mathcal{N}_{X, \sigma_c^+}) \cong H^3(\mathcal{N}_{X, \sigma_c^-})$ as required.
- If $d_1 - 2d_2 = 5$ then $\sigma_c = \sigma_m + \frac{3}{2} = \sigma_M - 6$. For $\sigma \in (\sigma_M - 6, \sigma_M - 3)$, we have $\mathcal{N}_{X, \sigma} = \mathcal{N}_{X, \sigma_L^-}$ for which the cohomology is computed above. For $\sigma \in (\sigma_m, \sigma_M - 6)$, we have that $\mathcal{N}_{X, \sigma} = \mathcal{N}_{X, \sigma_m^+}$. As $\mu_1 - \mu_2 > 2$, so have that $H^3(\mathcal{N}_{X, \sigma_m^+}) \cong H^3(M_X(2, L_0))$, where $L_0 = L_1 \otimes L_2^{-2}$ (see the proof of Theorem 5.5, and note that $\deg(L_0)$ is odd). We can twist L_0 by a large power of a line bundle μ to arrange that $\deg(L_0 \otimes \mu^{2k})$ is large. As $M_X(2, L_0) \cong M_X(2, L_0 \otimes \mu^{2k})$, we get that $H^3(\mathcal{N}_{X, \sigma_m^+}(2, 1, L_1, L_2))$ is isomorphic to $H^3(\mathcal{N}_{X, \sigma_m^+}(2, 1, L_1 \otimes \mu^{2k}, L_2))$. We have seen already that such cohomology group is isomorphic to $H^1(X)$ for $d_1 - 2d_2 \gg 0$. So $H^3(\mathcal{N}_{X, \sigma_c^-}) \cong H^3(\mathcal{N}_{X, \sigma_c^+})$ in this case.

The argument fails exactly for the critical value $\sigma_c = \sigma_M - 6 = \sigma_m + \frac{3}{2}$. However, Remark 5.2 implies that $H^3(\mathcal{N}_{X, \sigma_c}^s)$ is a mixed Hodge structure whose Gr_W^3 -piece is isomorphic to $H^1(X)$.

□

Now we shall compute the cohomology groups of $\mathcal{N}_{X, \sigma}^s$ and $M_X^s(n, L_0)$ simultaneously. We will prove the following two theorems.

Theorem 5.5. *Suppose $n \geq 2$, and $(n, g, d) \neq (2, 2, \text{even})$. Then*

- $H^1(M_X^s(n, L_0)) = 0$,
- $H^2(M_X^s(n, L_0)) \cong \mathbb{Z}$,
- $H^3(M_X^s(n, L_0)) \cong H^1(X)$, *except in the cases $(n, g, d) = (3, 2, 3k)$ and $(n, g, d) = (2, 3, 2k)$, $k \in \mathbb{Z}$. In these cases, $H^3(M_X^s(n, L_0))$ is a mixed Hodge structure and $\text{Gr}_W^3 H^3(M_X^s(n, L_0)) \cong H^1(X)$.*

(Recall that when n and d are coprime, $M_X(n, L_0) = M_X^s(n, L_0)$.)

Theorem 5.6. *Assume $n \geq 2$. Let $\sigma \in (\sigma_m, \sigma_M)$ if $n \geq 3$ and $\sigma \in (\sigma_m, \sigma_M - 3)$ if $n = 2$. Then*

- $H^1(\mathcal{N}_{X,\sigma}^s) = 0$,
- $H^2(\mathcal{N}_{X,\sigma}^s) \cong \mathbb{Z} \oplus \mathbb{Z}$,
- $H^3(\mathcal{N}_{X,\sigma}^s) \cong H^1(X)$,

except for the cases $(n, g, d_1 - 2d_2) = (2, 2, 5)$, $\sigma = \sigma_M - 6 = 4$, and $(n, g, d_1 - 3d_2) = (3, 2, 2k)$, $k = 1, 2, 3$.

We prove both Theorem 5.5 and 5.6 as follows. First we know that Theorem 5.6 is true for $n = 2$ by Proposition 5.4. Then we prove Theorem 5.5 for rank n assuming Theorem 5.6 for the same rank n . Finally we prove Theorem 5.6 for rank $n \geq 3$ using Theorem 5.5 for rank $n - 1$.

Proof of Theorem 5.5. Since the moduli spaces $M_X(n, L_0)$ and $M_X(n, L_0 \otimes \mu^n)$ are isomorphic via $E \mapsto E \otimes \mu$, for any fixed line bundle μ , we may assume that the degree d is large, say $d > (2g - 2)n$.

Fix $d_1 = d$, $d_2 = 0$, $L_1 = L_0$ and $L_2 = \mathcal{O}$, and consider the moduli spaces $\mathcal{N}_{X,\sigma} = \mathcal{N}_X(\sigma; n, 1, L_0, \mathcal{O})$. The moduli space $\mathcal{N}_{X,\sigma}$ for the smallest possible values of the parameter can be explicitly described. Let $\sigma_m^+ = \sigma_m + \epsilon$, $\epsilon > 0$ small enough. By [12, Proposition 4.10], there is a morphism

$$\pi : \mathcal{N}_{X,\sigma_m^+} \rightarrow M_X(n, L_0)$$

which sends $T = (E, L, \phi) \mapsto E$. Let $U = \pi^{-1}(M_X^s(n, L_0))$. By [12, Proposition 4.10], $\pi : U \rightarrow M_X^s(n, L_0)$ is a projective fibration whose fibers are the projective spaces $\mathbb{P}H^0(E)$, since $d_1/n - d_2 > 2g - 2$. Therefore

$$\begin{aligned} H^1(U) &\cong H^1(M_X^s(n, L_0)), \\ H^2(U) &\cong H^2(M_X^s(n, L_0)) \oplus \mathbb{Z}, \\ H^3(U) &\cong H^3(M_X^s(n, L_0)). \end{aligned}$$

Let us compute the cohomology groups of U . The complement $S = \mathcal{N}_{X,\sigma_m^+} - U$ consists of triples (E, L_2, ϕ) where E is semistable. By Lemma 4.3, the codimension of the family of such bundles is at least $(n - 1)(g - 1)$. The fiber over E is contained

in (but it may not be equal to) $\mathbb{P}H^0(E)$. As E is semistable and $d_1/n - d_2 > 2g - 2$, this dimension is constant. So $\text{codim } S \geq (n - 1)(g - 1)$.

If $(n, g, d) \neq (3, 2, 3k)$ and $(n, g, d) \neq (2, 3, 2k)$, $k \in \mathbb{Z}$, then $\text{codim } S \geq 3$. Then Lemma 5.1 implies that $H^1(U) = H^1(\mathcal{N}_{X, \sigma_m^+})$, $H^2(U) = H^2(\mathcal{N}_{X, \sigma_m^+})$ and $H^3(U) = H^3(\mathcal{N}_{X, \sigma_m^+})$. The result now follows from Theorem 5.6 for rank n .

If either $(n, g, d) = (3, 2, 3k)$ or $(n, g, d) = (2, 3, 2k)$, $k \in \mathbb{Z}$, then $\text{codim } S = 2$ and we only know by Remark 5.2 that $H^3(U)$ is a mixed Hodge structure with $\text{Gr}_W^3 H^3(U) = H^3(\mathcal{N}_{X, \sigma_m^+})$. $\square$

Proof of Theorem 5.6. We assume $n \geq 3$ since the case $n = 2$ is covered by Proposition 5.4.

Using Proposition 5.3, we see that it is enough to prove Proposition 5.7. $\square$

Proposition 5.7. *Assume $n \geq 3$, and assume that Theorem 5.5 holds for rank $n - 1$. Let $\sigma_M^- = \sigma_M - \epsilon$, $\epsilon > 0$ small enough. Then*

- $H^1(\mathcal{N}_{X, \sigma_M^-}) = 0$,
- $H^2(\mathcal{N}_{X, \sigma_M^-}) \cong \mathbb{Z} \oplus \mathbb{Z}$,
- $H^3(\mathcal{N}_{X, \sigma_M^-}) \cong H^1(X)$,

except for the case $(n, g, d_1 - 3d_2) = (3, 2, 2k)$, $k = 1, 2, 3$.

Proof. By Propositions 7.5 and 7.6 in [6], the triples in $\mathcal{N}_{X, \sigma_M^-}$ satisfy that $\phi : L_2 \rightarrow E_1$ is injective with torsion-free cokernel. Let $F = E_1/\phi(L_2)$. Then there is a short exact sequence $L_2 \rightarrow E_1 \rightarrow F$, and F is a semistable bundle. Moreover, the determinant of F is fixed, since $\det(F) = \det(E_1) \otimes L_2^{-1} = L_1 \otimes L_2^{-1}$. The extension is always non-trivial. Moreover, if F is stable, then any non-zero extension gives rise to a σ_M^- -stable triple.

By [6, Proposition 6.9], the dimension $\dim H^1(F^* \otimes L_2)$ is constant, for F as above. Let $U \subset \mathcal{N}_{X, \sigma_M^-}$ be the open subset formed by those triples with F a stable bundle. Then there is a fibration

$$U \rightarrow M_X^s(n - 1, L_1 \otimes L_2^{-1})$$

whose fibers are projective spaces $\mathbb{P}H^1(F^* \otimes L_2)$. Therefore

$$\begin{aligned} H^1(U) &= 0, \\ H^2(U) &\cong H^2(M_X^s(n - 1, L_1 \otimes L_2^{-1})) \oplus \mathbb{Z} \cong \mathbb{Z} \oplus \mathbb{Z}, \\ H^3(U) &\cong H^3(M_X^s(n - 1, L_1 \otimes L_2^{-1})). \end{aligned}$$

As the dimension $\dim H^1(F^* \otimes L_2)$ is constant, the codimension of $\mathcal{N}_{X, \sigma_M^-} - U$ is at least the codimension of a locus of semistable bundles. By Lemma 4.3 applied to $M_X(n - 1, L_1 \otimes L_2^{-1})$, this is at least $(n - 2)(g - 1)$. If $(n - 1, g, d_1 - d_2) \neq (2, 2, 2k)$, $(3, 2, 3k)$, $(2, 3, 2k)$, then this codimension is at least three. So $H^i(\mathcal{N}_{X, \sigma_M^-}) \cong H^i(U)$, for $i \leq 3$. Applying Theorem 5.5 for rank $n - 1$ we get the result.

If $(n-1, g, d_1-d_2) = (3, 2, 3k)$ or $(2, 3, 2k)$, then $\text{codim}(\mathcal{N}_{X, \sigma_M^-} - U) = 2$, so $H^3(\mathcal{N}_{X, \sigma_M^-}) \cong \text{Gr}_W^3 H^3(U)$. But again Theorem 5.5 gives us the result.

Suppose finally that $(n, g, d_1-d_2) = (3, 2, 2k)$, $k \in \mathbb{Z}$. By Proposition 5.3, $H^3(\mathcal{N}_{X, \sigma_M^-}) \cong H^3(\mathcal{N}_{X, \sigma_m^+})$. By the proof of Theorem 5.5,

$$H^3(\mathcal{N}_{X, \sigma_m^+}) \cong \text{Gr}_W^3 H^3(M_X^s(3, L_0)),$$

for $d_1/3 - d_2 > 2g - 2 = 2$, $L_0 = L_1 \otimes L_2^{-3}$ (note that we are not assuming that the right hand side is known). So assume $d_1 - 3d_2 > 6$. Twist L_0 by a line bundle μ of degree 1 so to change $\deg(L_0)$ to $\deg(L_0 \otimes \mu^{3k}) = \deg(L_0) + 3k$. This allows to change the parity of $d_1 - 3d_2$. Therefore $H^3(\mathcal{N}_{X, \sigma_m^+})$ is independent of the parity of $d_1 - d_2 \equiv d_1 - 3d_2 \pmod{2}$. Since the case that $d_1 - d_2$ is odd is already known, the result follows. $\square$

6. RECONSTRUCTING THE POLARISATION

We want to show that $H^3(M_X^s(n, L_0))$ and $H^3(\mathcal{N}_{X, \sigma}^s)$ have natural polarisations, which make them into polarised Hodge structures. The word “natural” means that they are constructed in families.

Proposition 6.1. *Suppose $(n, g, d) \neq (2, 2, 2k), (3, 2, 3k), (2, 3, 2k)$. The Hodge structure $H^3(M_X^s(n, L_0))$ is naturally polarised, and the isomorphism $H^3(M_X^s(n, L_0)) \cong H^1(X)$ respects the polarisations.*

Proof. For $M = M_X^s(n, L_0)$ the polarisation is constructed as follows (see [1, Section 8]; a similar argument is in [3, Section 4]). Let $\overline{M} = M_X(n, L_0)$. Since $H^2(M) = \mathbb{Z}$, we have that $\text{Pic}(M) = \mathbb{Z}$, so there is a unique ample generator of the Picard group. Take a general $(k-3)$ -fold hyperplane section $Z \subset M$, where $k = \dim M$. By Lemma 4.4, $\text{codim}(M_X(n, L_0) - M_X^s(n, L_0)) \geq 4$, so Z is smooth. Define

$$\begin{aligned} H^3(M) \otimes H^3(M) &\longrightarrow \mathbb{Z}, \\ \beta_1 \otimes \beta_2 &\mapsto \langle \beta_1 \cup \beta_2, [Z] \rangle. \end{aligned} \tag{6.1}$$

This is a polarisation. This is proved as follows (see [1, Proposition 6.2.1]): take a generic $(k-4)$ -fold hyperplane section $W \subset M$. As $\text{codim}(M_X(n, L_0) - M_X^s(n, L_0)) \geq 5$, W is smooth. By the Lefschetz theorem [1, Theorem 6.1.1] applied to the open smooth variety M , we have that $H^3(W) \cong H^3(M)$. Then by hard Lefschetz, cupping with the hyperplane class gives an isomorphism $H^3(W) \xrightarrow{\cong} H^5(W) \cong H^3(W)^*$. This map coincides with (6.1), which proves that it is a non-degenerate pairing.

Let $\pi : \mathcal{X} \rightarrow \mathcal{T}$ be the family of curves of genus g with no automorphisms, and let θ be the standard polarisation on $R^1\pi_*\mathbb{Z}$ corresponding to the cup product. We consider the universal Jacobian

$$q : \mathcal{J}^d \rightarrow \mathcal{T},$$

that is, the family of Jacobians $\text{Jac}^d X$, for $X \in \mathcal{T}$. Then there is a universal moduli space

$$p : \mathcal{M}_{\mathcal{T}}(n, d) \rightarrow \mathcal{J}^d$$

which puts over any $(X, L_0) \in \mathcal{J}^d$ the moduli space $M_X^s(n, L_0)$. If $\underline{\mathbb{Z}}$ denotes the local system over $\mathcal{M}_{\mathcal{T}}(n, d)$, then $R^3 p_* \underline{\mathbb{Z}}$ is the local system over $\mathcal{J}^d$ whose fibers are the Hodge structures $H^3(M_X^s(n, L_0))$. Let θ' be the natural polarisation on $R^3 p_* \underline{\mathbb{Z}}$ as defined above. There is an isomorphism

$$R^3 p_* \underline{\mathbb{Z}} \cong q^* R^1 \pi_* \underline{\mathbb{Z}}.$$

The natural map $\mathcal{J}^d \rightarrow M_g$ is dominant. So [1, Lemma 8.1.1] implies that there exists an integer $m \neq 0$ such that $\theta' = m\theta$. As any polarisation is a unique positive multiple of a primitive polarisation, θ' determines a unique primitive polarisation on $H^3(M_X^s(n, L_0))$ for any (X, L_0) . So the isomorphism $H^3(M_X^s(n, L_0)) \cong H^1(X)$ respects the polarisations. $\square$

Proposition 6.2. *Let $n \geq 2$. Let $\sigma \in (\sigma_m, \sigma_M)$ if $n \geq 3$ and $\sigma \in (\sigma_m, \sigma_M - 3)$ if $n = 2$. Assume that we are not in any of the cases enumerated in Lemma 4.2, and also that $(n, g, d_1 - 3d_2) \neq (3, 2, 2k)$, $k = 1, 2, 3$. Then $H^3(\mathcal{N}_{X,\sigma}^s)$ is naturally polarised, and the isomorphism $H^3(\mathcal{N}_{X,\sigma}^s) \cong H^1(X)$ respects the polarisations.*

Proof. Let $N = \mathcal{N}_{X,\sigma}^s = \mathcal{N}_X^s(\sigma; n, 1, L_1, L_2)$ and $\overline{N} = \mathcal{N}_{X,\sigma} = \mathcal{N}_X(\sigma; n, 1, L_1, L_2)$. By (2.4), we can assume $L_1 = L_0$ and $L_2 = \mathcal{O}$, with $d = \deg(L_0) = d_1 - nd_2$. As $H^2(N) \cong \mathbb{Z} \oplus \mathbb{Z}$, we have that $\text{Pic}(N) \cong \mathbb{Z} \oplus \mathbb{Z}$. Fix a basis H_1, H_2 for $\text{Pic}(N)$. For σ rational, $\overline{N}$ is naturally polarised, that is, there are $a, b \in \mathbb{Z}$ such that $H = aH_1 + bH_2$ is a (primitive) polarisation of N . Take a generic $(k-3)$ -fold hyperplane intersection $Z \subset N$. For σ non-critical, N is projective and smooth, so Z is smooth. For σ critical, Z is smooth as the codimension of the singular locus $\overline{N} - N$ is at least 4. Now consider the polarisation

$$\begin{aligned} H^3(N) \otimes H^3(N) &\longrightarrow \mathbb{Z}, \\ \beta_1 \otimes \beta_2 &\mapsto \langle \beta_1 \cup \beta_2, [Z] \rangle. \end{aligned}$$

This is a polarisation since $\text{codim}(\overline{N} - N) \geq 5$, which is proved as in Proposition 6.1.

Again let $\pi : \mathcal{X} \rightarrow \mathcal{T}$ be the family of curves of genus g with no automorphisms, and consider the universal Jacobian $q : \mathcal{J}^d \rightarrow \mathcal{T}$. Then there is a universal moduli space

$$p : \mathcal{N}_{\mathcal{T},\sigma} = \mathcal{N}_{\mathcal{T}}(\sigma; n, 1, d, \mathcal{O}) \rightarrow \mathcal{J}^d$$

which puts over any $(X, L_0) \in \mathcal{J}^d$ the moduli space $\mathcal{N}_X^s(\sigma; n, 1, L, \mathcal{O})$. There is a map $\mathcal{N}_{\mathcal{T},\sigma_M^-} \rightarrow \mathcal{M}_{\mathcal{T}}(n-1, d)$ defined on an open subset whose complement has codimension at least two. Pulling back the relative ample generator of $\mathcal{M}_{\mathcal{T}}(n-1, d) \rightarrow \mathcal{J}^d$, we get an element H_2 well defined in the family, $H_2 \in \text{Pic}(N)$. As p

is a projective bundle (off a subset of codimension at least two), there is another element H_1 which is well defined in the family, $H_1 \in \text{Pic}(N)$. The construction of the flips can be done in families, so $\text{Pic}(N) \cong \mathbb{Z}[H_1] \oplus \mathbb{Z}[H_2]$ with H_1, H_2 defined in families.

Let σ be non-critical. Then $H + \epsilon_1 H_1 + \epsilon_2 H_2$ is also a polarisation, for small rational ϵ_1, ϵ_2 . It is well-defined globally for the family. So the result [1, Lemma 8.1.1] gives us a rational number $m = m(\epsilon)$ so that

$$\beta_1 \cup \beta_2 \cup (H + \epsilon_1 H_1 + \epsilon_2 H_2)^{k-3} = m \theta(\beta_1, \beta_2) \quad \forall \beta_1, \beta_2 \in H^3(N).$$

This implies that $\beta_1 \cup \beta_2 \cup H_1^a \cup H_2^{k-3-a} = m_a \theta$, for some $m_a \in \mathbb{Q}$, for any $0 \leq a \leq k-3$. The conclusion is that for all possible polarisations θ' of N defined in families, we get that θ' is a multiple of θ .

If σ is critical, then if there is only one polarisation for $\mathcal{N}_{X,\sigma}^s$, there is nothing to prove. If there are several, then the ample cone contains an open set. So we can work as above to prove that all of the possible polarisations of $\mathcal{N}_{X,\sigma}^s$ give the same polarisation (up to multiples) for $H^3(\mathcal{N}_{X,\sigma}^s)$. $\square$

7. THE CASE OF NON-FIXED DETERMINANT

In this section we shall prove the Torelli theorem for the moduli spaces of pairs and bundles with non-fixed determinant, that is, Corollary 1.5.

We shall use the following lemma.

Lemma 7.1. *Let X be a projective connected variety and $f : X \rightarrow Y$ a map to another (quasi-projective) variety such that $f^* : H^k(Y) \rightarrow H^k(X)$ is zero for all $k > 0$. Then f is constant.*

Proof. Substituting X by an irreducible component, we can assume X is irreducible. Note that if f is constant on each irreducible component, then it is constant, by the connectedness of X .

Let d be the dimension of a generic fiber of f , and consider a generic d -fold hyperplane intersection $Z \subset X$, which is transverse to the generic fiber. Then $f|_Z : Z \rightarrow Y$ is a proper and generically finite map. Therefore $f|_Z : Z \rightarrow f(Z)$ is of finite degree $N > 0$, and $f(Z) \subset Y$ is a closed subvariety. Therefore $H^{2t}(f(Z)) \cong \mathbb{Z} \rightarrow H^{2t}(Z) \cong \mathbb{Z}$, $t = \dim Z$, is multiplication by N . If $t > 0$, the assumption of the lemma implies that $H^{2t}(Y) \rightarrow H^{2t}(f(Z))$ should be zero. But this is impossible, since a generic t -fold hyperplane intersection in Y maps to a non-zero element in $H^{2t}(f(Z))$. Therefore $t = 0$, i.e. f is constant. $\square$

Proof of Corollary 1.5. Let $\mathfrak{M}_X(\tau; n, d)$ be the moduli space of τ -polystable pairs of rank n and degree d . There is a determinant map

$$\det : \mathfrak{M}_X(\tau; n, d) \rightarrow \text{Jac}^d X,$$

sending $(E, \phi) \mapsto \det(E)$, whose fiber over L_0 is the moduli space $\mathfrak{M}_X(n, L_0)$.

Assume that $F : \mathfrak{M}_X(\tau; n, d) \xrightarrow{\cong} \mathfrak{M}_{X'}(\tau'; n', d')$, for X' another curve. Fix a line bundle L'_0 on X' of degree d' and consider the composition

$$f : \mathfrak{M}_{X'}(\tau'; n', L'_0) \hookrightarrow \mathfrak{M}_{X'}(\tau'; n', d') \cong \mathfrak{M}_X(\tau; n, d) \rightarrow \text{Jac}^d X.$$

As $f^* : H^1(\text{Jac}^d X) \rightarrow H^1(\mathfrak{M}_{X'}(\tau'; n, L_0)) = 0$ is the zero map, and $H^*(\text{Jac}^d X)$ is generated by $H^1(\text{Jac}^d X)$, we have that the map $f^* : H^k(\text{Jac}^d X) \rightarrow H^k(\mathfrak{M}_{X'}(\tau'; n, L_0))$ is zero for all $k > 0$. Applying Lemma 7.1, we have that f is constant. Therefore there exists a line bundle L_0 on X of degree d such that F maps $M' = \mathfrak{M}_{X'}(\tau'; n', L'_0)$ to $M = \mathfrak{M}_X(\tau; n, L_0)$. Working analogously, the map F^{-1} maps $\mathfrak{M}_X(\tau; n, L_0)$ into some fiber of the map $\det$, which must be M' . This implies that $F|_{M'} : M' \rightarrow M$ is an isomorphism. Now we apply Corollary 1.2 to conclude that $X \cong X'$.

Suppose that τ' is a critical value and that there is an isomorphism $F : \mathfrak{M}_X^s(\tau; n, d) \cong \mathfrak{M}_{X'}^s(\tau'; n', d')$. Then we have a map

$$f : M' = \mathfrak{M}_{X'}^s(\tau'; n', L'_0) \hookrightarrow \mathfrak{M}_{X'}^s(\tau'; n', d') \cong \mathfrak{M}_X^s(\tau; n, d) \rightarrow \text{Jac}^d X.$$

Now take any compactification $\bar{M}'$ of M' . So there is a rational map $f : \bar{M}' \dashrightarrow \text{Jac}^d X$. After blowing-up, we have a compactification $\tilde{M}'$ of M' and a map $\tilde{f} : \tilde{M}' \rightarrow \text{Jac}^d X$ which extends f . As $H^1(M') = 0$, we have that $H^1(\tilde{M}') = 0$ as well. Using Lemma 7.1, we have that $\tilde{f}$ is a constant map. The rest of the argument is as before.

The case of bundles is entirely analogous. □

REFERENCES

- [1] ARAPURA, D.; SASTRY, P.: Intermediate Jacobians and Hodge structures of moduli spaces. *Proc. Indian Acad. Sci. Math. Sci.* **110** (2000) 1–26.
- [2] BERTRAM, A.: Stable pairs and stable parabolic pairs. *J. Algebraic Geom.* **3** (1994), no. 4, 703–724.
- [3] BISWAS, I.; MUÑOZ, V.: The Torelli theorem for the moduli spaces of connections on a Riemann surface. *Topology* **46** (2007), no. 3, 295–317.
- [4] BRADLOW, S. B.; DASKALOPOULOS: Moduli of stable pairs for holomorphic bundles over Riemann surfaces. *Internat. J. Math.* **2** (1991) 477–513.
- [5] BRADLOW, S. B.; GARCÍA-PRADA, O.: Stable triples, equivariant bundles and dimensional reduction. *Math. Ann.* **304** (1996) 225–252.
- [6] BRADLOW, S. B.; GARCÍA-PRADA, O.; GOTHEN, P.B: Moduli spaces of holomorphic triples over compact Riemann surfaces. *Math. Ann.* **328** (2004) 299–351.
- [7] BRADLOW, S. B.; GARCÍA-PRADA, O.; MUÑOZ, V.; NEWSTEAD, P. E.: Coherent systems and Brill-Noether theory. *Internat. J. Math.* **14** (2003), no. 7, 683–733.
- [8] GARCÍA-PRADA, O.: Dimensional reduction of stable bundles, vortices and stable pairs. *Internat. J. Math.* **5** (1994) 1–52.
- [9] KOUVIDAKIS, A.; PANTEV, T.: The automorphism group of the moduli space of semistable vector bundles. *Math. Ann.* **302** (1995) 225–268.

- [10] MUMFORD, D.; NEWSTEAD, P.: Periods of a moduli space of bundles on curves. *Amer. J. Math.* **90** (1968) 1200–1208.
- [11] MUÑOZ, V.: Hodge polynomials of the moduli spaces of rank 3 pairs. *Geometriae Dedicata*. To appear.
- [12] MUÑOZ, V.; ORTEGA, D.; VÁZQUEZ-GALLO, M.-J.: Hodge polynomials of the moduli spaces of pairs. *Internat. J. Math.* **18** (2007) 695–721.
- [13] NARASIMHAN, M. S.; RAMANAN, S.: Geometry of Hecke cycles. I. C. P. Ramanujam—a tribute, pp. 291–345, *Tata Inst. Fund. Res. Studies in Math.* **8**, Springer, Berlin-New York, 1978.
- [14] NARASIMHAN, M. S.; RAMANAN, S.: Moduli of vector bundles on a compact Riemann surface, *Ann. of Math. (2)* **89** (1969) 14–51.
- [15] SCHMITT, A.: A universal construction for the moduli spaces of decorated vector bundles. *Transform. Groups* **9** (2004) 167–209.
- [16] THADDEUS, M.: Stable pairs, linear systems and the Verlinde formula. *Invent. Math.* **117** (1994) 317–353.
- [17] TYURIN, A. N.: Geometry of moduli of vector bundles. *Russ. Math. Surveys* **29** (1974) 59–88.

INSTITUTO DE CIENCIAS MATEMÁTICAS CSIC-UAM-UCM-UC3M, CONSEJO SUPERIOR DE INVESTIGACIONES CIENTÍFICAS, SERRANO 113 BIS, 28006 MADRID, SPAIN

FACULTAD DE MATEMÁTICAS, UNIVERSIDAD COMPLUTENSE DE MADRID, PLAZA CIENCIAS 3, 28040 MADRID, SPAIN

E-mail address: vicente.munoz@imaff.cfmac.csic.es